FORM PTO 1390

3.

US LE PARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. §371

ATTORNEY DOCKET NUMBER 2001\_1838A

U.S. APPLICATION NO. 962

International Application No. PCT/JP00/03806

International Filing Date
June 12, 2000

**Priority Date Claimed** June 14, 1999

#### **Title of Invention**

PLANT THERMOGENIC GENES AND PROTEINS

#### Applicant(s) For DO/EO/US

Kikukatsu ITO

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

- 1. [X] This is a FIRST submission of items concerning a filing under 35 U.S.C. §371.
- 2. [] This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. §371.
- 3. [] This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1).
- 4. [X] A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
- 5. [X] A copy of the International Application as filed (35 U.S.C. §371(c)(2))
  - a. [] is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. [X] has been transmitted by the International Bureau.
  - c. [] is not required, as the application was filed in the United States Receiving Office (RO/US)
- 6. [X] A translation of the International Application into English (35 U.S.C. §371(c)(2)). ATTACHMENT A
- 7. [] Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3)).
  - a. [] are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. [] have been transmitted by the International Bureau.
  - c. [] have not been made; however, the time limit for making such amendments has NOT expired.
  - d. [] have not been made and will not be made.
- 8. [] A translation of the amendments to the claims under PCT Article 19.
- 9. [X] An unexecuted oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)). ATTACHMENT B
- 10. [] A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)).

#### Items 11. to 14. below concern other document(s) or information included:

- 11. [X] An Information Disclosure Statement under 37 CFR 1.97 and 1.98. ATTACHMENT C
- 12. [] An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
- 13. [] A FIRST preliminary amendment.
  - [] A SECOND or SUBSEQUENT preliminary amendment.
- 14. [] Other items or information:

THE COMMISSIONER IS AUTHORIZED TO CHARGE ANY DEFICIENCY IN THE FEE FOR THIS PAPER TO DEPOSIT ACCOUNT NO. 23-0975.

TO THE REAL PROPERTY OF THE PR									
U.S. APPLICATION NO. (4 1000, 1000) 962 INTERNATIONAL APPLICATION NO. PCT/JP00/03806				ATTORNEY'S DOCKET NO. 2001_1838A					
15. [X] The following fees are su	CALCULATIONS	PTO USE ONLY							
BASIC NATIONAL FE Neither international preliminary and International Search Report International Search Report has b International preliminary examina paid to USPTO International preliminary examina of PCT Article 33(1)-(4) International preliminary examina PCT Article 33(1)-(4)									
ENTER APPROI	\$890.00								
Surcharge of \$130.00 for furnishic claimed priority date (37 CFR 1.4	\$								
Claims	Number Filed	Number Extra	Rate						
Total Claims	-20 =		X \$18.00	\$					
Independent Claims	4 - 3 =	1	X \$84.00	\$84.00					
Multiple dependent claim(s) (if ap	plicable)		+ \$280.00	\$					
TOTAL	\$974.00								
Small Entity Status is here	\$								
	\$974.00								
Processing fee of \$130,00 for furn claimed priority date (37 CFR 1.4	\$	-							
	\$974.00								
Fee for recording the enclosed ass appropriate cover sheet (37 CFR 3	\$	-							
	\$974.00								
	Amount to be refunded	s							
······································	Amount to be charged	\$							
<ul> <li>a. [X] A check in the amount of \$974.00 to cover the above fees is enclosed. A duplicate copy of this form is enclosed.</li> <li>b. [] Please charge my Deposit Account No. 23-0975 in the amount of \$ to cover the above fees.  A duplicate copy of this sheet is enclosed.</li> </ul>									
c. [] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-0975.									
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.									
19. CORRESPONDENCE ADDR	9. CORRESPONDENCE ADDRESS								
	0513		By: Warren M. Cheek, Jr., Registration No. 33,367  WENDEROTH, LIND & PONACK, L.L.P. 2033 "K" Street, N.W., Suite 800  Washington, D.C. 20006-1021  Phone: (202) 721-8200						
	202) 721-8250								

47962 [CHECK NO. [2001\_1838A]

December 14, 2001

## JC09 Ree'd PCT/PTO

1 4 DEC 2001 ATTORNEY DOCKET NUMBER PORM PTO 1390 (REV 3-93) US DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE 2001 1838A TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) U.S. APPLICATION NO. CONCERNING A FILING UNDER 35 U.S.C. §371 **Priority Date Claimed International Filing Date** International Application No. June 14, 1999 June 12, 2000 PCT/JP00/03806 Title of Invention PLANT THERMOGENIC GENES AND PROTEINS Applicant(s) For DO/EO/US Kikukatsu ITO Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: 1. [X] This is a FIRST submission of items concerning a filing under 35 U.S.C. §371. 2. [] This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. §371. 3. [] This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1). 4. [X] A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. [X] A copy of the International Application as filed (35 U.S.C. §371(c)(2)) a. [] is transmitted herewith (required only if not transmitted by the International Bureau). b. [X] has been transmitted by the International Bureau. c. [] is not required, as the application was filed in the United States Receiving Office (RO/US) 6. [X] A translation of the International Application into English (35 U.S.C. §371(c)(2)). ATTACHMENT A 7. [] Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3)). a. [] are transmitted herewith (required only if not transmitted by the International Bureau). b. [] have been transmitted by the International Bureau. c. [] have not been made; however, the time limit for making such amendments has NOT expired. d. [] have not been made and will not be made. 8. [] A translation of the amendments to the claims under PCT Article 19. 9. [X] An unexecuted oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)). ATTACHMENT B 10. [] A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)). Items 11. to 14. below concern other document(s) or information included: 14. [X] An Information Disclosure Statement under 37 CFR 1.97 and 1.98. ATTACHMENT C 12. [] An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.

> THE COMMISSIONER IS AUTHORIZED TO CHARGE ANY DEFICIENCY IN THE FEE FOR THIS PAPER TO DEPOSIT **ACCOUNT NO. 23-0975.**

13. [] A FIRST preliminary amendment.

14. [] Other items or information:

[] A SECOND or SUBSEQUENT preliminary amendment.

U.S. APPLICATION NO.			TION NO.	ATTORNEY'S DOCKET NO. 2001 1838A				
15. [X] The following fees are submitted					CALCULATIONS	PTO USE ONLY		
BASIC NATIONAL FE Neither international preliminary and International Search Report International Search Report has b International preliminary examina paid to USPTO International preliminary examina of PCT Article 33(1)-(4) International preliminary examina PCT Article 33(1)-(4)								
ENTER APPRO	\$890.00							
Surcharge of \$130.00 for furnish claimed priority date (37 CFR 1.	\$	,						
Claims	Number Filed	Number I	Extra	Rate				
Total Claims	-20 =			X \$18.00	s			
Independent Claims	4 - 3 =	1		X \$84.00	\$84.00			
Multiple dependent claim(s) (if a	\$							
TOTAL OF ABOVE CALCULATIONS =					\$974.00			
[] Small Entity Status is here	\$	·						
	\$974.00	-						
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).					\$			
TOTAL NATIONAL FEE =					\$974.00	•,		
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40 per property +					\$	Ç		
	\$974.00							
					Amount to be refunded	\$		
					Amount to be charged	\$		
a. [X] A check in the amount of \$974.00 to cover the above fees is enclosed. A duplicate copy of this form is enclosed.								
b. [] Please charge my Deposit Account No. 23-0975 in the amount of \$ to cover the above fees.  A duplicate copy of this sheet is enclosed.								
c. [] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-0975.								
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.								
19. CORRESPONDENCE ADD	RESS			11	1. 11. 11	2		

000513
PATENT TRADEMARK OFFICE

By:

Warren M. Cheek, Jr., Registration No. 33,367

WENDEROTH, LIND & PONACK, L.L.P. 2033 "K" Street, N.W., Suite 800 Washington, D.C. 20006-1021 Phone:(202) 721-8200 Fax:(202) 721-8250

December 14, 2001

[CHECK NO. 47962

[2001\_1838A]



# Recopes: 012842

#7/

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Confirmation No. 3213

Kikukatsu ITO

Docket No. 2001-1838A

Serial No. 10/009,962

Group Art Unit Not Yet Assigned

Filed January 23, 2002

Examiner Not Yet Assigned

PLANT THERMOGENIC GENES AND PROTEINS

TO CHARGE ANY DEFICIENCY IN THE FES FOR THIS PAPER TO DEPOSIT ACCOUNT NO. 23-0975

## **PRELIMINARY AMENDMENT**

Assistant Commissioner for Patents, Washington, D.C. 20231

Sir:

Responsive to the Notice dated February 13, 2002, the time for filing thereto being extended for one month in accordance with the Petition for Extension submitted concurrently herewith, please amend the above-identified application as follows:

### In the Specification:

Page 1, line 1, delete the entire heading.

between lines 3 and 6, insert the following new heading:

Background of the Invention

line 6, replace the heading with the following new heading:

1. Field of the Invention

line 16, replace the heading with the following new heading:

2. Description of the Related Art

## Page 3, replace the paragraph beginning at line 11 with the following paragraph:

Potato and *Arabidopsis* have been considered to be non-thermogenic plants.

However, the expression of StUCP and AtPUMP have been induced by low temperature.

Therefore, it has been suggested that these genes are involved in heat production (Laloi et al., 1997; Maia et al., 1998).

line 26, replace the heading with the following new heading: Summary of the Invention

## Page 4, replace the paragraph beginning at line 23 with the following paragraph:

Fig. 3 compares the alignment of amino acid sequences of SfUCPA (SEQ ID No. 2) and SfUCPB (SEQ ID No. 4), together with potato UCP (StUCP) (SEQ ID No. 5), Arabidopsis UCP (AtPUMP) (SEQ ID No. 6) and human UCP (human UCP 1, 2 and 3 corresponding to SEQ ID Nos. 7, 8 and 9, respectively). The asterisks (\*) attached under the sequences indicate the same amino acid sequence, and the dot (.) indicates the conservative change in all of the sequences. The boldface indicates the same sequence between SfUCPA (SEQ ID No. 2) and SfUCPB (SEQ ID No. 4). The gap introduced to optimize the sequence alignment is indicated by a dash (-). The alignment was made using a CLUSTAL W program. The characteristic domains of energy transfer proteins typical of mitochondria are surrounded by a square. The shaded bars (1~VI) above the upper sequence show estimated transmembrane domains.

Page 5, line 25, replace the heading with the following new heading:

Description of the Preferred Embodiments

## Page 10, replace the paragraph beginning at line 28 with the following paragraph:

- 10mM Tris-HC1 (pH 8.0);
- 50mM KC1;
- 1.5mM MgC1<sub>2</sub>;
- · 4mM dNTP;
- 0.2 unit of EX Taq polymerase (Takara); and
- 10pmol of two degenerate primers corresponding to the conserved amino acid sequence of the UCP family:

ZF1 (5'-CCIYTIGAYACIGCIAAR-3') (SEQ ID No. 11)

ZR1 (5'-ACWTTCCAISYICCIAWIC-3') (SEQ ID No. 12).

Page 13, line 15, delete the entire heading.

## In the Claims:

Page 16, above claim 1, insert the following:

What is claimed is:

#### In the Abstract:

Page 17, line 1, replace the heading with the following new heading

ABSTRACT OF THE DISCLOSURE

## In the Sequence Listing:

Please replace the Sequence Listing of record with the attached substitute Sequence Listing consisting of SEQ ID Nos. 1-12.

#### **REMARKS**

The foregoing amendments are presented to place the application in compliance with the sequence rules under 37 CFR 1.821-1.825.

Applicants have submitted a Sequence Listing in both paper and computer readable form as required by 37 C.F.R. 1.821(c) and (e). Amendments directing its entry into the specification have also been incorporated herein. The content of the paper and computer readable copies are the same and no new matter has been added.

Additional amendments to the specification have also been effected to put the specification in better form under U.S. practice. Specifically, the specification headings have been amended in

THE CARLEST CONTRACTOR

conformance with U.S. practice. Also, the additional sequences disclosed in the specification and

Figure 3 have been identified and labeled as required under U.S. practice.

With regard to the Notice also requesting that an executed Oath and Declaration of the

Inventors needs to be submitted, Applicants wish to note that an executed Oath and Declaration

was submitted on January 23, 2002. A copy of the submitted executed Declaration is enclosed

herewith along with the cover letter (indicating the filing of the executed Declaration). Applicants

respectfully request that the Patent Office review the application papers to ensure that the

executed Declaration is present in the file.

Attached hereto is a marked-up version of the changes made to the specification by the

current amendment. The attached page is captioned "Version with markings to show changes

made."

In view of the foregoing, it is believed that each requirement set forth in the Notice has

been satisfied, and that the application is now in compliance with the sequence rules under 37

CFR 1.821-1.825. Accordingly, favorable examination on the merits is respectfully requested.

Respectfully submitted,

Kikukatsu ITO

By: Lee Cheng

Registration No. 40,949

Attorney for Applicant

LC/gtn


Washington, D.C. 20006-1021

Telephone (202) 721-8200

Facsimile (202) 721-8250

May 13, 2002

- 5 -



5

10

20

25

#### DESCRIPTION

1

#### Plant Thermogenic Genes and Proteins

Background of the Invention InTechnical Field of the Invention

The present invention relates to plant thermogenic genes and proteins. More particularly, the invention relates to thermpogenic genes derived from a skunk cabbage (Symplocarpus foetidus) and gene products (proteins). Those genes and proteins are useful in breeding of cold-avoidance plants, medical treatment of diabetes mellitus or obesity, or development of novel thermogenic bio-materials.

Stresses due to low temperatures, droughts and salinity are common harmful environmental factors that terrestrial plants encounter. Among these stresses, it has been considered that cellular injury due to the low temperature is the most important factor which restricts productivity of crops (Levitt, 1980). To resist the low temperature stress, cold-hardy plants such as wheat or rye have a variety of physiological and metabolic responses which lead to cold acclimation (Sakai and Larcher, 1987; Steponkus, 1984; Thomashow, 1998; Uemura and Steponkus, 1997). In contrast, it is known that some plants including skunk cabbage have a specialized system by which the plants generate heat to avoid freezing (Knutson, 1974; Nagy et al., 1972; Schneider and Buchenen, 1980).

The temperature of the flower in the spadix of skunk cabbage, which 30 flowers in early spring, has been known to maintain its temperature at higher the C-terminal region and increases by free fatty acids (Jezek et al., 1998; Lin and Klingenberg, 1982; Katiyar and Shrago, 1989; Rial et al., 1983; Sluse et al., 1998).

On the contrary, 2 cDNAs encoding UCP-like proteins of plant origin were isolated from potato (StUCP: Laloi et al., 1997) and from *Arabidopsis* (AtPUMP: Maia et al., 1998). Since the expression of StUCP was mainly detected in the flower and the fruit, it has been postulated that StUCP may concern respiration during flowering and maturation of the fruit together with the AOX activity (Laloi et al., 1997).

10

5

Potato and Arabidopsis have been considered to be non-thermogenic plants. However, the expression of StUCP and AtPUMP induced by low temperature. Therefore, it has been suggested that these genes are involved in the heat production (Laloi et al., 1997; Maia et al., 1998).

15

In the thermogenic plants such as skunk cabbage, however, UCP-mediated thermogenic mechanisms have not yet been identified.

The purpose of the invention of this application is to provide unidentified novel UCP genes derived from a thermogenic plant, skunk cabbage.

The additional purpose of this application is to provide skunk cabbage UCPs which are expression products of the novel genes.

25

30

20

# Summary of the Disclosure of Invention

The invention provides thermogenic genes derived from skunk cabbage, i.e., gene SfUCPa of which cDNA comprises the base sequence of SEQ ID NO: 1, and gene SfUCPb of which cDNA comprises the base sequence of SEQ ID NO: 3.

Moreover, the invention provides thermogenic proteins, i.e., protein SfUCPA expressed from SfUCPa, which comprises the amino acid sequence of SEQ ID NO: 2, and protein SIUCPB expressed from SIUCPb, which comprises the amino acid sequence of SEQ ID NO: 4.

In addition, the invention provides cDNA having the base sequence of SEQ ID NO: 1 or a partial sequence thereof, and cDNA having the base sequence of SEO ID NO: 3 or a partial sequence thereof.

10

20

25

5

## **Brief Description of Drawings**

Fig. 1 shows the change of the temperature of the spadix in skunk cabbage and that of ambient temperature with a lapse of time. 15

Fig. 2 shows the results of northern blotting, indicating the expression profile of SfUCPa (A) and SfUCPb (B) in the spadix and leaf of skunk cabbage at room temperature (RT) and during cold treatment (4°C for 3 days). The lower figures respectively show the results of ethidium bromide staining of non-decomposed rRNA.

(SEQ ID No.2 Fig. 3 compares the alignment of amino acid sequences of SfUCPA and (SEQ ID No. 4) (SEQ ID NO. 5) (SEQ ID NO. 5) (SEQ ID NO. 5) (SEQ ID NO. 5) (SEQ ID NO. 5)

human UCP. The asterisks (\*) attached under the sequences indicate the same (human UCP), 2 and 3 correspond to SEQ JD Nbs. 7, 8 and 9, respectively amino acid sequence, and the dot (.) indicates the conservative change in all of the The boldface indicates the same sequence between SfUCPA and sequences. SIUCPB. The gap introduced to optimize the sequence alignment is indicated by a dash (-). The alignment was made using a CLUSTAL W program. characteristic domains of energy transfer proteins typical of mitochondria are 30

surrounded by a square. The shaded bars (I~VI) above the upper sequence show estimated transmembrane domains.

Fig. 4 shows a hydrophobic plot of SfUCPA. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM6.

Fig. 5 shows a diagrammatic illustration of SfUCPA topology in the mitochondria membrane.

10

15

Fig. 6 shows a hydrophobic plot of SfUCPB. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM4 and TM6.

Fig. 7 shows a diagrammatic illustration of SfUCPB topology in the mitochondria membrane.

Fig. 8 shows the results of *in vitro* translation using respective cDNAs of the genes SfUCPa and SfUCPb as templates. (-) indicates a control, S a sense RNA, and AS an antisense RNA. The asterisk (\*) indicates a non-specific product and the empty circle denotes the position of a low molecular translated artificial product synthesized from a small ORF.

# Description of the Preferred Embodiments Best Mode for Carrying Out the Invention

25

30

20

In the gene SfUCPa of the present invention, its cDNA has the base sequence of SEQ ID NO: 1 and encodes the protein SfUCPA having the amino acid sequence of SEQ ID NO: 2, of which the estimated molecular weight is 32.6 kDa. In the gene SfUCPb of the present invention, its cDNA (SEQ ID NO: 3) encodes

many include mammal cultured cells such as monkey renal cell COS7, Chinese hamster ovarian cell CHO, etc., budding yeast, fission yeast, silkworm cell, Xenopus egg cell, and the like are commonly used, but not limited thereto. In order to introduce the expression vector into eucaryotic cells, a known method such as electroporation, calcium phosphate method, liposome method, DEAE dextran method, and the like can be utilized.

After expression of the proteins in procaryotic cells or eucaryotic cells according to the aforementioned method, the desired proteins are isolated and purified from the culture in the known combined procedures for separation. For example, treatment with a denaturant (e.g., urea) or surface activator, ultrasonication, digestion with enzymes, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity chromatography, reverse phase chromatography, and the like are involved.

The proteins of the invention, SfUCPA and SfUCPB, also include peptide fragments (5 amino acids or more) involving the optional partial amino acid sequences of SEQ ID NOS: 2 and 4. In addition, the proteins of the invention also include fusion proteins with other optional proteins.

The following examples serve to illustrate the invention of this application specifically in more detail, but are not intended to limit the scope of the invention.

25

30

5

10

15

20

#### Example 1: Cloning of cDNA

The total RNA was extracted from the spadix of skunk cabbage (Symplocarpus foetidus) and the complete RNA was determined on 1.0% agarose gel electrophoresis (Ito et al., 1999). Using a mRNA isolation kit (Pharmacia), a

- 10mM Tris-HCl (pH 8.0);
- 50mM KCl;
- 10 1.5mM MgCl<sub>2</sub>;
  - 4mM dNTP;
  - 0.2 unit of EX Taq polymerase (Takara); and
  - 10pmol of two degenerate primers corresponding to the conserved amino acid sequence of the UCP family:

ZF1 (5'-CCIYTIGAYACIGCIAAR-3') (SEQ ID NO II)
ZR1 (5'-ACWTICCAISYICCIAWIC-3')(SEQ ID NO IO)

PCR cycle was carried out as follows.

(94°C: 0.5 minute; 50°C: 1 minute; 72°C: 1 minute)×35

20

25

30

15

5

Among the PCR products obtained in the above method, the amino acid sequence estimated from the sequence of about 0.8kb cDNA fragment had very high homology to one of the reading frame sequences of the UCP gene family. This fragment, accordingly, was cloned into T-vector (clone p2-1) and used as a probe for library screening.

cDNA (5µg) prepared from the spadix was inserted into  $\lambda gt11$  phage according to the known method (Sambrook et al., 1989) to construct a cDNA library. From this library, 8 clones positive to the above-described probes were isolated and sub-cloned into the pBluescript SK plasmid (Stratagene). From

linearized, on which a sense- or anti-sense RNA was transcribed with T7 RNA polymerase or T3 RNA polymerase according to the protocol of MAXICRIPT transcription kit (Ambion). An equal amount of RNA (4µg) was provided for in vitro translation reaction using a wheat germ extract (Promega) in the presence of <sup>35</sup>S-methionine (Amersham). The translation product was analyzed by SDS-PAGE. The gel was fixed, incubated in Amplify (Amersham), then dried, and fluorometrically analyzed.

As a result, it was confirmed that, as shown in Fig. 8, the initiation codon and the stop codon of cDNA isolated in Example 1 functioned successfully since a protein having an expected molecular weight was produced from any of cDNAs only when the sense RNA was used as a template.

#### Industrial Applicability

As described previously, this application provides novel thermogenic genes SfUCPa and SfUCPb as well as their gene products, i.e., thermogenic proteins SfUCPA and SfUCPB, derived from skunk cabbage (Symplocarpus foetidus), and cDNAs used for gene engineering mass production of these proteins. These genes and proteins allow development of low temperature-tolerant plants, development of drugs or methods for treatment of diabetes mellitus or obesity, or development of novel heat generating materials from plants.

25

20

5

10

15

#### References

Berthold and Siedow (1993) Plant Physiol. 101, 113-119. Boss et al. (1997) FEBS Lett. 408, 39-42.

30 Fleury et al. (1997) Nature Genetics 15, 269-272.

#### **CLAIMS**

What is daimed is !

- 1. A thermogenic gene SfUCPa derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 1.
- 2. A thermogenic gene SfUCPb derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 3.
- 3. A thermogenic protein SfUCPA expressed from the gene SfUCPa of Claim
  10 1, which comprises the amino acid sequence of SEQ ID NO: 2.
  - 4. A Thermogenic protein SfUCPB expressed from the gene SfUCPb of Claim 2, which comprises the amino acid sequence of SEQ ID NO: 4.
- 5. A DNA fragment comprising the base sequence of SEQ ID NO: 1 or a partial sequence thereof.
  - 6. A DNA fragment comprising the base sequence of SEQ ID NO 3: or a partial sequence thereof.

20

5

## ABSTRACT OF THE DISCLOSULE

The inventions of this application include thermogenic genes named SfUCPa and SfUCPb which are derived from skunk cabbage. cDNA of each gene comprises the base sequence of SEQ ID NO: 1 and 3, respectively. Thermogenic proteins, SfUCPA and SfUCPB, are expressed from genes SfUCPa and SfUCPb, comprises the amino acid sequence of SEQ ID NO: 2 and 4.

5

B/PRTS

JC05 Reva PGT/PTO 1 4 DEC 2001

#### DESCRIPTION

#### Plant Thermogenic Genes and Proteins

5

10

#### **Technical Field**

The present invention relates to plant thermogenic genes and proteins. More particularly, the invention relates to thermpogenic genes derived from a skunk cabbage (*Symplocarpus foetidus*) and gene products (proteins). Those genes and proteins are useful in breeding of cold-avoidance plants, medical treatment of diabetes mellitus or obesity, or development of novel thermogenic bio-materials.

15

20

25

30

#### Background

Stresses due to low temperatures, droughts and salinity are common harmful environmental factors that terrestrial plants encounter. Among these stresses, it has been considered that cellular injury due to the low temperature is the most important factor which restricts productivity of crops (Levitt, 1980). To resist the low temperature stress, cold-hardy plants such as wheat or rye have a variety of physiological and metabolic responses which lead to cold acclimation (Sakai and Larcher, 1987; Steponkus, 1984; Thomashow, 1998; Uemura and Steponkus, 1997). In contrast, it is known that some plants including skunk cabbage have a specialized system by which the plants generate heat to avoid freezing (Knutson, 1974; Nagy et al., 1972; Schneider and Buchenen, 1980).

The temperature of the flower in the spadix of skunk cabbage, which flowers in early spring, has been known to maintain its temperature at higher

than +10°C even when the ambient temperature falls to -15°C (Knutson, 1974). For example, thermoscopic analysis using infrared camera indicates homeothermic behavior of the surface temperature of the spadix (Fig. 1). It should be noted that, in this experiment, the plants were placed in the growth chamber and the air temperature was gradually decreased. As clearly seen from Fig. 1, the temperature of the spadix of skunk cabbage is kept at approximately 19°C notwithstanding a fall of the ambient temperature.

The temperature is thus maintained by doubling the respiration rate from the level of 12°C to that of sub-zero temperature. It has also been considered that the heat production in thermogenic plant species relates to a cellular metabolism called cyanide-non-sensitive/non-phosphorylating electron-transferring pathway, which is controlled by mitochondrial alternative oxidase (AOX)(Berthold and Siedow, 1993; Ito et al., 1997; McIntosh, 1994, Wangner and Krab, 1995).

On the other hand, it has been shown that a mitochondrial protein called an uncoupling protein (UCP) plays an important role in generation of heat in mammals. UCP found in the intima of mitochondria make H+ flow into the membrane to uncouple aspiration from synthesis of ATP which acts to disperse chemical energy to metabolic heat (Klaus et al., 1991; Klingenberg and Winkler, 1985; Ricquier et al., 1991). In animals, 3 types of UCPs have been found. UCP1 is primarily distributed in brown adipose tissue (Nichollus and Locke, 1984). UCP2 is found ubiquitously in many tissues (Fleury et al., 1997), and UCP3 is localized specifically in skeletal muscle (Boss et al., 1997).

It has been considered that UCPs of mammals, similarly to other carrier proteins of mitochondria, are composed of 6 transmembrane segments, of which the hydrophobic portion is derived from pairing amphipathic  $\alpha$ -helix structure (Liu et al., 1988; Maia et al., 1998). It is also known that the activity of these UCPs decreases depending on purine nucleotides (ATP, GTP, GDP and ADP) attached to

the C-terminal region and increases by free fatty acids (Jezek et al., 1998; Lin and Klingenberg, 1982; Katiyar and Shrago, 1989; Rial et al., 1983; Sluse et al., 1998).

On the contrary, 2 cDNAs encoding UCP-like proteins of plant origin were isolated from potato (StUCP: Laloi et al., 1997) and from *Arabidopsis* (AtPUMP: Maia et al., 1998). Since the expression of StUCP was mainly detected in the flower and the fruit, it has been postulated that StUCP may concern respiration during flowering and maturation of the fruit together with the AOX activity (Laloi et al., 1997).

10

5

Potato and *Arabidopsis* have been considered to be non-thermogenic plants. However, the expression of StUCP and AtPUMP induced by low temperature. Therefore, it has been suggested that these genes are involved in the heat production (Laloi et al., 1997; Maia et al., 1998).

15

In the thermogenic plants such as skunk cabbage, however, UCP-mediated thermoigenic mechanisms have not yet been identified.

The purpose of the invention of this application is to provide unidentified novel UCP genes derived from a thermogenic plant, skunk cabbage.

The additional purpose of this application is to provide skunk cabbage UCPs which are expression products of the novel genes.

25

30

#### Disclosure of Invention

The invention provides thermogenic genes derived from skunk cabbage, i.e., gene SfUCPa of which cDNA comprises the base sequence of SEQ ID NO: 1, and gene SfUCPb of which cDNA comprises the base sequence of SEQ ID NO: 3.

Moreover, the invention provides thermogenic proteins, i.e., protein SfUCPA expressed from SfUCPa, which comprises the amino acid sequence of SEQ ID NO: 2, and protein SfUCPB expressed from SfUCPb, which comprises the amino acid sequence of SEQ ID NO: 4.

In addition, the invention provides cDNA having the base sequence of SEQ ID NO: 1 or a partial sequence thereof, and cDNA having the base sequence of SEQ ID NO: 3 or a partial sequence thereof.

10

20

25

30

5

#### **Brief Description of Drawings**

Fig. 1 shows the change of the temperature of the spadix in skunk cabbage and that of ambient temperature with a lapse of time.

Fig. 2 shows the results of northern blotting, indicating the expression profile of SfUCPa (A) and SfUCPb (B) in the spadix and leaf of skunk cabbage at room temperature (RT) and during cold treatment (4°C for 3 days). The lower figures respectively show the results of ethidium bromide staining of non-decomposed rRNA.

Fig. 3 compares the alignment of amino acid sequences of SfUCPA and SfUCPB, together with potato UCP (StUCP), *Arabidopsis* UCP (AtPUMP) and human UCP. The asterisks (\*) attached under the sequences indicate the same amino acid sequence, and the dot (.) indicates the conservative change in all of the sequences. The boldface indicates the same sequence between SfUCPA and SfUCPB. The gap introduced to optimize the sequence alignment is indicated by a dash (-). The alignment was made using a CLUSTAL W program. The characteristic domains of energy transfer proteins typical of mitochondria are

surrounded by a square. The shaded bars (I~VI) above the upper sequence show estimated transmembrane domains.

Fig. 4 shows a hydrophobic plot of SfUCPA. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM6.

Fig. 5 shows a diagrammatic illustration of SfUCPA topology in the mitochondria membrane.

10

15

20

25

30

Fig. 6 shows a hydrophobic plot of SfUCPB. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM4 and TM6.

Fig. 7 shows a diagrammatic illustration of SfUCPB topology in the mitochondria membrane.

Fig. 8 shows the results of *in vitro* translation using respective cDNAs of the genes SfUCPa and SfUCPb as templates. (-) indicates a control, S a sense RNA, and AS an antisense RNA. The asterisk (\*) indicates a non-specific product and the empty circle denotes the position of a low molecular translated artificial product synthesized from a small ORF.

#### Best Mode for Carrying Out the Invention

In the gene SfUCPa of the present invention, its cDNA has the base sequence of SEQ ID NO: 1 and encodes the protein SfUCPA having the amino acid sequence of SEQ ID NO: 2, of which the estimated molecular weight is 32.6 kDa. In the gene SfUCPb of the present invention, its cDNA (SEQ ID NO: 3) encodes

the protein SfUCPB having the amino acid sequence of SEQ ID NO: 4, of which the estimated molecular weight is 29.0 kDa.

Genes SfUCPa and SfUCPb of the invention are derived from skunk cabbage, which are expressed specifically in the spadix when the temperature is low. The results of Northern blotting on the total RNAs extracted from skunk cabbage (Ito et al., 1999), confirmed that the expressions of both genes were detected in the spadices but not in the leaves at room temperature (15°C) (Fig. 2). It was also confirmed that the spadix-specific expression of both genes were induced by cold treatment (4°C for 3 days).

5

10

15

20

25

30

The amino acid sequences of the proteins SfUCPA and SfUCPB that are expressed from the respective genes of the invention have higher homology to the plant UCPs than to the human UCPs (Fig. 3) such that the amino acid sequence of SfUCPA has homology of 79%, 75%, 44%, 48% and 48% to StUCP, AtPUMP, human UCP, UCP2 and UCP3, respectively. SfUCPB has homology of 71%, 66%, 41%, 43% and 44% to StUCP, AtPUMP, human UCP, UCP2 and UCP3, respectively.

In addition, SfUCPA and SfUCPB have high sequence homology (88%) to each other though the region corresponding to the amino acid sequence between the 204th Thr and the 238th Val in SfUCPA is completely deleted in SfUCPB (Fig. 3). Moreover, the 265th Leu of SfUCPA is replaced by Pro in SfUCPB.

StUCPA has similar structure to that of other mitochondria UCP proteins. SfUCPA has 6 transmembrane domains as shown by the hydrophobic plot in Fig. 4, of which the topology is as shown in Fig. 5. In addition, this SfUCPA has 3 domains that are characteristic of energy transfer proteins in mitochondria (Fig. 3)(Boss et al., 1997; Maia et al., 1998). On the other hand, SfUCPB is lacking in the 3rd domain which is characteristic of energy transfer proteins in mitochondrial

(Fig. 3), as well as in the 5th transmembrane domain (Figs. 3 and 6). The topology is located toward the mitochondria matrix at the C-terminal (Fig. 7).

Each protein has a purine nucleotide-binding domain (PNBD) at the C-terminal (Figs. 3, 5 and 7), and it is known that in UCP, binding of the purine nucleotide inhibits the uncoupling function in the mitochondria intima. In SfUCPB, however, there is a possibility that it may have escaped the inhibition of the binding of the purine nucleotide because its C-terminal is located toward the mitochondria matrix. Such a topology has not been found in any UCPs from animals or plants.

5

10

15

20

25

30

The thermogenic genes SfUCPa and SfUCPb provided by the invention are derived from skunk cabbage and are very useful in, for example, development of low temperature-tolerant plants using a genetic recombination technique. The proteins SfUCPA and SfUCPB that are expression products from the above genes are expected as effective components in remedies of diabetes mellitus, obesty, and the like, based on the uncoupling function to ATP synthesis. Moreover, such thermogenic proteins are also promising novel heat generating bio-materials.

The genes SfUCPa and SfUCPb of the invention can be isolated from the genomic DNA of skunk cabbage using the cDNA (SEQ ID NOS: 1 or 3) or a partial sequence thereof of the invention as a probe. For example, a genome library is prepared from the genomic DNA according to a known method. It may be screened by means of colony or plaque hybridization according to a known method using as a probe an oligonucleotide synthesized based on the base sequence of an optional portion of cDNA. Alternatively, the target genetic region may also be identified by means of in situ hybridization for chromosome.

The respective cDNAs of the invention can be cloned, for example, from a cDNA library which is synthesized using a poly(A)+RNA of skunk cabbage as a

template. In such a case, an oligonucleotide of an optional portion of cDNA provided by the invention is synthesized, which may be used as a probe to carry out screening by means of colony or plaque hybridization according to a known method. Alternatively, oligonucleotides which can hybridize to both ends of the target cDNA fragment are synthesized, which may be used as primers in preparation of cDNA of the invention by the RT-PCR method from mRNA isolated from the cells of skunk cabbage.

5

20

25

30

In general, polymorphism is frequently recognized in the genes of eucaryotic cells. In the invention, accordingly, in addition to cDNAs represented by SEQ ID NOS: 1 and 3, those in which one or several nucleotides are added, deleted and/or replaced by (an)other nucleotide(s) in the above cDNA are included. Similarly, proteins in which one or more amino acids are added, deleted and/or replaced by (an)other amino acid(s) due to change of the above nucleotide are also included in the present invention.

In cDNA of the invention, DNA fragments (10bp or more) comprising an optimal part of the base sequences of SEQ ID NOS 1 and 3 are included. In addition, DNA fragments comprising a sense strand or anti-sense strand are also included.

The proteins of the invention, SfUCPA and SfUCPB, may be prepared respectively by a known method, for example, isolation from the spadix of skunk cabbage, preparation by chemical syntheses based on the amino acid sequence provided by the invention, or production by a recombinant DNA technique using cDNA provided by the invention. For example, when the protein is produced by a recombinant DNA technique, RNA is prepared from a vector containing cDNA of the invention by *in vitro* transcription, and this is used as a template for *in vitro* translation to yield the protein. Alternatively, the translational region of cDNA is incorporated into an appropriate expression vector according to a known method,

and the resulting recombinant vector is introduced into *Escherichia coli*, *Baccillus subtilis*, yeast, animal or plant cells. The resulting transformants can be used in expression of the proteins in a large quantity.

In the case of the proteins of the invention being produced by *in vitro* translation, the translation region of cDNA of the invention may be incorporated into a vector containing RNA polymerase promotor, and then added to an *in vitro* translation system such as a rabbit reticulocyte lysate or wheat germ extract containing an RNA polymerase corresponding to the promotor. The RNA polymerase promotor is exemplified by T7, T3, SP6, and similar promoters.

In the case of the proteins of the invention being expressed in a microorganism such as *Escherichia coli*, the translation region of cDNA is incorporated into an expression vector containing an origin replicable in microorganisms, promoter, ribosome binding site, cDNA cloning site, terminator, and the like, to construct a recombinant expression vector, which is then introduced into a host cell and incubated. In this operation, an initiation codon and a stop codon may be added to the front and tail of an optional translation region to obtain a protein fragment containing the optional region. Alternatively, the desired protein may be expressed as a fusion protein with another protein, which may be cleaved with a suitable protease to yield the desired protein. The expression vectors for *Escherichia coli* are exemplified by pUC series, pBluescript II, pET expression system, pGEX expression system, and the like.

In the case of the proteins of the invention being expressed in eucaryotic cells, the translation region of cDNA of the invention is incorporated into an expression vector for eucaryotic cells containing a promoter, splicing region, poly(A) additional site, and the like, and introduced into the eucaryotic cells. The expression vector is exemplified by pKA1, pCDM8, pSVK3, pMSG, pSVL, pBK-CMV, pBK-RSV, EVB-vector, pRS, pYES2, and the like. The eucaryotic cells,

many include mammal cultured cells such as monkey renal cell COS7, Chinese hamster ovarian cell CHO, etc., budding yeast, fission yeast, silkworm cell, Xenopus egg cell, and the like are commonly used, but not limited thereto. In order to introduce the expression vector into eucaryotic cells, a known method such as electroporation, calcium phosphate method, liposome method, DEAE dextran method, and the like can be utilized.

After expression of the proteins in procaryotic cells or eucaryotic cells according to the aforementioned method, the desired proteins are isolated and purified from the culture in the known combined procedures for separation. For example, treatment with a denaturant (e.g., urea) or surface activator, ultrasonication, digestion with enzymes, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity chromatography, reverse phase chromatography, and the like are involved.

The proteins of the invention, SfUCPA and SfUCPB, also include peptide fragments (5 amino acids or more) involving the optional partial amino acid sequences of SEQ ID NOS: 2 and 4. In addition, the proteins of the invention also include fusion proteins with other optional proteins.

The following examples serve to illustrate the invention of this application specifically in more detail, but are not intended to limit the scope of the invention.

## Example 1: Cloning of cDNA

5

10

15

20

25

30

The total RNA was extracted from the spadix of skunk cabbage (Symplocarpus foetidus) and the complete RNA was determined on 1.0% agarose gel electrophoresis (Ito et al., 1999). Using a mRNA isolation kit (Pharmacia), a

- 10mM Tris-HCl (pH 8.0);
- 50mM KCl;
- 1.5mM MgCl<sub>2</sub>;
  - 4mM dNTP;
  - 0.2 unit of EX Taq polymerase (Takara); and
  - 10pmol of two degenerate primers corresponding to the conserved amino acid sequence of the UCP family:

ZF1 (5'-CCIYTIGAYACIGCIAAR-3')

ZR1 (5'-ACWTTCCAISYICCIAWIC-3')

PCR cycle was carried out as follows.

(94°C: 0.5 minute; 50°C: 1 minute; 72°C: 1 minute)×35

20

25

30

5

10

15

Among the PCR products obtained in the above method, the amino acid sequence estimated from the sequence of about 0.8kb cDNA fragment had very high homology to one of the reading frame sequences of the UCP gene family. This fragment, accordingly, was cloned into T-vector (clone p2-1) and used as a probe for library screening.

cDNA (5 $\mu$ g) prepared from the spadix was inserted into  $\lambda$ gt11 phage according to the known method (Sambrook et al., 1989) to construct a cDNA library. From this library, 8 clones positive to the above-described probes were isolated and sub-cloned into the pBluescript SK plasmid (Stratagene). From

these clones, clones pz8-1 and pz8-2 were obtained, which respectively contained the full length SfUCPa cDNA and SfUCPb cDNA.

The insert in each clone was sequenced with an auto-sequencer ABI373A using the BcaBest sequencing kit (Takara) and T3, T7 and gene-specific primers. The sequence data were analyzed by means of the GENETYX-Homology Software System version 2.2.0 (Software Development).

cDNA of SfUCPa had the 1,525bp base sequence of SEQ ID NO: 1, and cDNA of SfUCPb had the 2,991bp base sequence of SEQ ID NO: 3. An estimated polyadenylated signal (aataaa) was found upstream of 236bp from the poly(A) sequence in cDNA of SfUCPa, while in cDNA of SfUCPb two polyadenylated sites were recognized at the positions of 1,171bp and 1,243bp. It is noteworthy that cDNA of SfUCPb has a longer 3'-untranslation region than that of SfUCPa.

15

20

25

10

5

cDNA of SfUCPa had an open reading frame (ORF) encoding 303 amino acids as shown in SEQ ID NO: 1, and this ORF was found to encode the protein SfUCPA of the estimated molecular weight 32.6kDa having the amino acid sequence of SEQ ID NO: 2. On the other hand, cDNA of SfUCPb had an ORF corresponding to 268 amino acids as shown in SEQ ID NO: 3, and found to encode the protein SfUCPB of the estimated molecular weight 29.0kDa.

Moreover, it was confirmed from the results of Southern blot analysis that the genome of skunk cabbage contains multiple copies of SfUCPa gene and a single copy of SfUCPb (data not shown).

#### Example 2: In vitro translation of cDNA

The plasmid clones pz8-1 and pz8-2 obtained in Example 1 were

linearized, on which a sense- or anti-sense RNA was transcribed with T7 RNA polymerase or T3 RNA polymerase according to the protocol of MAXICRIPT transcription kit (Ambion). An equal amount of RNA (4µg) was provided for in vitro translation reaction using a wheat germ extract (Promega) in the presence of 35S-methionine (Amersham). The translation product was analyzed by SDS-PAGE. The gel was fixed, incubated in Amplify (Amersham), then dried, and fluorometrically analyzed.

As a result, it was confirmed that, as shown in Fig. 8, the initiation codon and the stop codon of cDNA isolated in Example 1 functioned successfully since a protein having an expected molecular weight was produced from any of cDNAs only when the sense RNA was used as a template.

#### **Industrial Applicability**

As described previously, this application provides novel thermogenic genes SfUCPa and SfUCPb as well as their gene products, i.e., thermogenic proteins SfUCPA and SfUCPB, derived from skunk cabbage (Symplocarpus foetidus), and cDNAs used for gene engineering mass production of these proteins. These genes and proteins allow development of low temperature-tolerant plants, development of drugs or methods for treatment of diabetes mellitus or obesity, or development of novel heat generating materials from plants.

25

20

5

10

15

#### References

Berthold and Siedow (1993) Plant Physiol. 101, 113-119. Boss et al. (1997) FEBS Lett. 408, 39-42.

30 Fleury et al. (1997) Nature Genetics 15, 269-272.

Ito, K. et al. (1999) Plant Sci. 142, 57-65.

Ito, K. et al. (1994) Nucl. Acids Res. 22, 2036-2041.

Ito, Y. et al. (1997) Gene 12, 121-129.

Jezek et al. (1998) Biochem. Biophys. Acta 1365, 319-327.

5 Katiyar and Shrago (1989) Nati. Acad. Sci. USA 86, 2559-2562.

Klaus et al. (1991) Int. J. Biochem. 23, 791-810.

Klingenberg and Winkler (1985) EMBO J. 4, 3087-3092.

Knutson (1974) Science 186, 746-747.

Laloi et al. (1997) Nature 389, 135-136.

10 Levitt (1980) Responses of plants to environmental stresses. 2nd edn. New York: Academic Press.

Lin and Klingenberg (1982) Biochemistry 21, 2950-2956.

Liu et al. (1988) J. Cell. Biol. 107, 503-509.

Maia et al. (1998) FEBS Lett. 429, 403-406.

15 McIntosh (1994) Plant Physiol. 105, 781-786.

Nagy et al. (1972) Science 178, 1195-1197.

Nicholls and Locke (1984) Physiol. Rev. 64, 1-64.

Rial et al. (1983) Eur. J. Biochem. 137, 197-203.

Ricquier et al. (1991) FASEB J. 5, 2237-2242.

20 Sakai and Larcher (1987) Frost Survival of Plants: Responses and Adaptations to Freezing Stresses. Berlin and New York; Springer-Verlag.

Sambrook et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd eds. New York; Cold Spring Harbor Laboratory Press.

Schneider and Buchanan (1980) Amer. J. Bot. 67, 182-193.

25 Sluse et al. (1998) FEBS Lett. 433, 237-240.

Steponkus (1984) Annu. Rev. Plant Physiol. 35, 543-581.

Thomashow (1998) Plant Physiol. 118, 1-7.

Uemura and Steponkus (1997) Plant Cold Hardiness: Molecular Biology, Biochemistry, and Physiology (Li and Chen, edn). New York: Plenum Press, pp.

30 171-179.

Wagner and Krab (1995) Physiol. Plant, 95, 318-325.

#### **CLAIMS**

- 1. A thermogenic gene SfUCPa derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 1.
- 2. A thermogenic gene SfUCPb derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 3.
- 3. A thermogenic protein SfUCPA expressed from the gene SfUCPa of Claim
  10. 1, which comprises the amino acid sequence of SEQ ID NO: 2.
  - 4. A Thermogenic protein SfUCPB expressed from the gene SfUCPb of Claim 2, which comprises the amino acid sequence of SEQ ID NO: 4.
- 5. A DNA fragment comprising the base sequence of SEQ ID NO: 1 or a partial sequence thereof.
  - 6. A DNA fragment comprising the base sequence of SEQ ID NO 3: or a partial sequence thereof.

5

#### **ABSTRACT**

The inventions of this application include thermogenic genes named SfUCPa and SfUCPb which are derived from skunk cabbage. cDNA of each gene comprises the base sequence of SEQ ID NO: 1 and 3, respectively. Thermogenic proteins, SfUCPA and SfUCPB, are expressed from genes SfUCPa and SfUCPb, comprises the amino acid sequence of SEQ ID NO: 2 and 4.

5

#### SEQUENCE LISTING

<110> Japan Science and Technology Corporation

<120> Plant Thermogenic Genes and Proteins

<150> JP11-167439

<151> 1999-06-14

<160> 4

<170> Patentin Ver. 2.0

<210> 1

<211> 1525

<212> DNA

<213> Symplocarpus foetidus

<220>

<221> CDS

<222> (280).. (1188)

<221> poly A site

<221> (1271).. (1276)

<300>

<301> I to, K.

<302> Isolation of two distinct cold-inducible cDNAs encoding plant uncoupling proteins from the spadix of skunk cabbage (Symplocarpus foetidus)

<303> Plant Sci.

<304> 149

<305>

<306> 167-173

<307> 1999

<308> GenBank AB024733

<309> 2000-02-25

<400> 1

gagg	atto	gc a	igaag	aaag	g cc	agaa	ccce	att	cctt	ССС	gtct	tctt	ct	ctto	cgccc	60
aatt	gcag	tt 1	ttce	cago	g gc	gtca	tcat	caa	gaco	ctc	cgcc	tttc	cg	gcca	aacgc	120
cttc	cacc	cc o	cacco	aato	g co	ctcc	gttt	ccc	gaaa	tat	tcct	ctto	cc t	ccto	ccttt	180
tctt	ctct	ac a	ataaa	ccct	a ac	caco	ccat	cct	ctc	tcc	cgct	tcce	gac o	acco	tgcat	240
tcta	ctgg	ga g	gccca	tttg	ga to	gage	gtttc	CCE	gcga	igg a	tg g	gc e	gat o	ac g	gc	294
										N	let 0	ily #	\sp ⊦	lis (	ìlу	
											1				5	
ccg	agg	acc	gag	atc	tcg	ttt	gcc	ggc	agt	tcg	cga	gca	gca	ttc	gcc	342
Pro	Arg	Thr	Glu	llе	Ser	Phe	Ala	Gly	Ser	Ser	Arg	Ala	Ala	Phe	Ala	
				10					15					20		
gct	tgc	ttc	gcc	gag	ctt	tgc	acg	att	ccg	ttg	gac	act	gct	aaa	gtt	390
Ala	Cys	Phe	Ala	Glu	Leu	Cys	Thr	Пe	Pro	Leu	Asp	Thr	Ala	Lys	Val	
			25					30					35			
agg	ctt	caa	ctc	caa	aag	aaa	gca	gta	aca	ggt	gat	gtg	gtg	gct	ttg	438
Arg	Leu	GIn	Leu	Gln	Lys	Lys	Ala	Val	Thr	Gly	Asp	Val	Val	Ala	Leu	
		40					45					50				
cca	aaa	tac	agg	gga	atg	ttg	ggc	act	gtt	gcc	act	att	gcc	agg	gag	486
Pro	Lys	Tyr	Arg	Gly	Met	Leu	Gly	Thr	Val	Ala	Thr	He	Ala	Arg	Glu	
	55					60					65					
gaa	ggt	ttg	tcg	gca	ctc	tgg	aaa	gga	att	gta	ccc	ggt	ttg	cat	cgt	534
Glu	Gly	Leu	Ser	Ala	Leu	Trp	Lys	Gly	He	Val	Pro	Gly	Leu	His	Arg	
70					75					80					85	
caa	tgc	ctc	ttt	gga	ggg	cta	cga	att	ggg	ttg	tat	gaa	cca	gtt	aag	582
Gln	Cys	Leu	Phe	Gly	Gly	Leu	Arg	He	Gly	Leu	Tyr	Glu	Pro	Val	Lys	
				90					95					100		
tcc	ttt	tat	gtt	gga	gat	aac	ttt	gtt	gga	gat	att	cct	tta	tcc	aag	630
Ser	Phe	Tyr	Val	Gly	Asp	Asn	Phe	Val	Gly	Asp	He	Pro	Leu	Ser	Lys	
			105					110					115			

aaa	ata	ctt	gct	ggg	ctt	aca	aca	ggt	gca	tta	gca	att	ata	gtg	gca	678
Lys	He	Leu	Ala	Gly	Leu	Thr	Thr	Gly	Ala	Leu	A I <sup>.</sup> a	Пe	He	Val	Ala	
		120					125					130				
aat	ссс	act	gac	ctt	gtt	aaa	gtt	cga	ctt	caa	tct	gaa	ggt	aaa	ctc	726
Asn	Pro	Thr	Asp	Leu	Val	Lys	Val	Arg	Leu	Gln	Ser	Glu	Gly	Lys	Leu	
	135					140					145					
CCC	cct	ggg	gta	ccg	aga	cgt	tat	tca	ggg	gcg	cta	aat	gct	tat	tca	774
Pro	Pro	Gly	Val	Pro	Arg	Arg	Tyr	Ser	Gly	Ala	Leu	Asn	Ala	Tyr	Ser	
150					155					160					165	
acc	ata	gtc	aaa	aag	gaa	gga	ctt	ggt	gct	ctg	tgg	act	ggg	ctt	ggt	822
Thr	He	Val	Lys	Lys	Glu	Gly	Leu	Gly	Ala	Leu	Trp	Thr	Gly	Leu	Gly	
				170					175					180		
cct	aat	att	gcc	cgc	aat	gct	att	ata	aat	gct	gct	gaa	ttg	gcc	agt	870
Pro	Asn	He	Ala	Arg	Asn	Ala	Пe	He	Asn	Ala	Ala	Glu	Leu	Ala	Ser	
			185					190					195			
tat	gat	caa	gtg	aaa	cag	aca	atc	tta	aaa	tta	cca	gga	ttc	tca	gat	918
Tyr	Asp	GIn	Val	Lys	GIn	Thr	lle	Leu	Lys	Leu	Pro	Gly	Phe	Ser	Asp	
		200					205					210				
aat	att	ttt	act	cat	att	tta	gcc	ggt	ctg	ggg	gca	ggt	ttt	ttt	gcc	966
Asn	He	Phe	Thr	His	lle	Leu	Ala	Gly	Leu	Gly	Ala	Gly	Phe	Phe	Ala	
	215					220					225					
gtc	tgt	atc	ggt	tct	cct	gtt	gat	gtg	atg	aag	tct	aga	atg	atg	gga	1014
Val	Cys	He	Gly	Ser	Pro	Val	Asp	Val	Met	Lys	Ser	Arg	Met	Met	Gly	
230					235					240					245	
gat	tca	gcc	tac	aaa	agc	aca	ttt	gat	tgt	ttc	atc	aag	aca	ttg	aaa	1062
Asp	Ser	Ala	Tyr	Lys	Ser	Thr	Phe	Asp	Cys	Phe	Пe	Lys	Thr	Leu	Lys	
				250					255					260		
22+	and t	aaa	c++	ctt	ac t	+++	tac	220	aaa	+++	ato	000	220	+++	aa t	1110

Asn Asp Gly Leu Leu Ala Phe Tyr Lys Gly Phe lle Pro Asn Phe Gly 265 270 275 cgg tta gga tcg tgg aat gtg atc atg ttt ttg acc ttg gag cag gtc 1158 Arg Leu Gly Ser Trp Asn Val IIe Met Phe Leu Thr Leu Glu Gin Val 280 285 290 aag aag tit itc atc aaa gag gig cca aat taatacatig aactcggata 1208 Lys Lys Phe Phe IIe Lys Glu Val Pro Asn 295 300 ggagtagaaa gaaagggttt ttgtggaatt ttctctaccg gtgtggatcc tggcgagaga 1268 caaataaatc ttcctgactg ctcagatgtg tacctttttt atgaatggtt cttttcttat 1328 agaggacaga gaaaagaaaa aaaaaatcat tgtcatttac tctttttccc catttctgct 1388 gctaatcttg gtaggagaag aaaagtctta cattgagtga taacgttgtt ctctgcatcc 1448 attatttttc agagatacta tttgacacat gaaaagtaat gcacatcagg ttttttttaa 1508 aaaaaaaaa aaaaaaa 1525 <210> 2 <211> 303 <212> PRT <213> Symplocarpus foetidus <400> 2 Met Gly Asp His Gly Pro Arg Thr Glu IIe Ser Phe Ala Gly Ser Ser 5 10 15 1 Arg Ala Ala Phe Ala Ala Cys Phe Ala Glu Leu Cys Thr lle Pro Leu 25 20 30 Asp Thr Ala Lys Val Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly 35 40 45 Asp Val Val Ala Leu Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala

55

60

50

Thr	He	Ala	Arg	Glu	Glu	Gly	Leu	Ser	Ala	Leu	Trp	Lys	Gly	He	Val
65					70					75					80
Pro	Gly	Leu	His	Arg	GIn	Cys	Leu	Phe	Gly	Gly	Leu	Arg	He	Gly	Leu
				85					90					95	
Tyr	Glu	Pro	Val	Lys	Ser	Phe	Tyr	Val	Gly	Asp	Asn	Phe	Val	Gly	Asp
			100					105					110		
He	Pro	Leu	Ser	Lys	Lys	He	Leu	Ala	Gly	Leu	Thr	Thr	Gly	Ala	Leu
		115					120					125			
Ala	He	Пe	Val	Ala	Asn	Pro	Thr	Asp	Leu	Val	Lys	Val	Arg	Leu	GIn
	130					135					140				
Ser	Glu	Gly	Lys	Leu	Pro	Pro	Gly	Val	Pro	Arg	Arg	Tyr	Ser	Gly	Ala
145					150					155					160
Leu	Asn	Ala	Tyr	Ser	Thr	lle	Val	Lys	Lys	Glu	Gly	Leu	Gly	Ala	Leu
				165					170					175	
Trp	Thr	Gly	Leu	Gly	Pro	Asn	He	Ala	Arg	Asn	Ala	Пe	lle	Asn	Ala
			180					185					190		
Ala	Glu	Leu	Ala	Ser	Tyr	Asp	Gln	Val	Lys	Gln	Thr	Пe	Leu	Lys	Leu
		195					200					205			
Pro	Gly	Phe	Ser	Asp	Asn	He	Phe	Thr	His	He	Leu	Ala	Gly	Leu	Gly
	210					215					220				
Ala	Gly	Phe	Phe	Ala	Val	Cys	He	Gly	Ser	Pro	Val	Asp	Val	Met	Lys
225					230					235					240
Ser	Arg	Met	Met	Gly	Asp	Ser	Ala	Tyr	Lys	Ser	Thr	Phe	Asp	Cys	Phe
				245					250					255	
He	Lys	Thr	Leu	Lys	Asn	Asp	Gly	Leu	Leu	Ala	Phe	Tyr	Lys	Gly	Phe
			260					265					270		
He	Pro	Asn	Phe	Gly	Arg	Leu	Gly	Ser	Trp	Asn	Val	He	Met	Phe	Leu
		275					280					285			

Thr Leu Glu Gln Val Lys Lys Phe Phe lle Lys Glu Val Pro Asn 290 295 300

<210> 3

<211> 2991

<212> DNA

<213> Symplocarpus foetidus

<220>

<221> CDS

<222> (286) . . (1089)

<221> poly A site

<222> (1171) (1176)

<221> poly A site

<222> (1243) (1248)

<300>

<301> I to, K.

<302> Isolation of two distinct cold-inducible cDNAs encoding plant uncoupling proteins from the spadix of skunk cabbage (Symplocarpus foetidus)

<303> Plant Sci.

<304> 149

<305>

<306> 167-173

<307> 1999

<308> GenBank AB024734

<309> 2000-02-25

<400> 3

tggtggtgac gagtgacgag gattcgcaga agaaaggcca gaacccgatt ccttcccgtc 60 ttcttctcct tccgcccaat tgcagttttt cgcagcgggt catcatcaag accctccgcc 120

tttccgcgcc aaacgccttc cacccaatcc ctccgtttcc cgaaatattc cccttccctc 180
ccttttcttc tctacataaa ccctaaccac ccccatcctc tcctcccgct tccgaccacc 240
ctgcattcta ctgggatccc atttgatcga cgtttcccgg cgagg atg ggc gat cac 297
Met Gly Asp His

													1			
ggc	ccg	agg	асс	gag	atc	tcg	ttt	gcc	ggc	agt	tcg	cga	gca	gca	ttc	345
Gly	Pro	Arg	Thr	Glu	Пе	Ser	Phe	Ala	Gly	Ser	Ser	Arg	Ala	Ala	Phe	
5					10					15					20	
gcc	gct	tgc	ttc	gcc	gag	ctc	tgt	acg	att	ccg	ttg	gac	act	gct	aaa	393
Ala	Ala	Cys	Phe	Ala	Glu	Leu	Cys	Thr	He	Pro	Leu	Asp	Thr	Ala	Lys	
				25					30					35		
gtt	agg	ctt	cag	ctc	caa	aag	aaa	gca	gta	aca	ggt	gat	gtg	gtg	gct	441
Val	Arg	Leu	Gln	Leu	Gln	Lys	Lys	Ala	Val	Thr	Gly	Asp	Val	Val	Ala	
			40					45					50			
ttg	cca	aaa	tac	agg	gga	atg	ttg	ggc	act	gtt	gcc	act	att	gcc	agg	489
Leu	Pro	Lys	Tyr	Arg	Gly	Met	Leu	Gly	Thr	Val	Ala	Thr	He	Ala	Arg	
		55					60					65				
gag	gaa	ggt	ttg	tcg	gca	ctc	tgg	aaa	gga	att	gta	ccc	ggt	ttg	cat	537
Glu	Glu	Gly	Leu	Ser	Ala	Leu	Trp	Lys	Gly	lle	Val	Pro	Gly	Leu	His	
	70					75					80					
cgt	caa	tgc	ctc	ttt	gga	ggg	cta	cga	att	ggg	ttg	tat	gaa	cca	gtt	585
Arg	GIn	Cys	Leu	Phe	Gly	Gly	Leu	Arg	He	Gly	Leu	Tyr	Glu	Pro	Val	
85					90					95					100	
aag	tcc	ttt	tat	gtt	gga	gat	aac	ttt	gtt	gga	gat	att	cct	tta	tcc	633
Lys	Ser	Phe	Tyr	Val	Gly	Asp	Asn	Phe	Val	Gly	Asp	Пe	Pro	Leu	Ser	
				105					110					115		
aag	aaa	ata	ctt	gct	ggg	ctt	aca	aca	ggt	gca	tta	gca	att	ata	gtg	681
Lys	Lys	He	Leu	Ala	Gly	Leu	Thr	Thr	Gly	Ala	Leu	Ala	He	He	Val	

			120					125					130			
gca	aat	ccg	act	gac	ctt	gtt	aaa	gtt	cga	ctt	caa	tct	gaa	ggt	aaa	729
Ala	Asn	Pro	Thr	Asp	Leu	Val	Lys	Val	Arg	Leu	Gln	Ser	Glu	Gly	Lys	
		135					140					145				
ctc	ссс	cct	ggg	gta	сса	aga	cgt	tat	tca	ggg	gcg	cta	aat	gct	tat	777
Leu	Pro	Pro	Gly	Val	Pro	Arg	Arg	Tyr	Ser	Gly	Ala	Leu	Asn	Ala	Tyr	
	150					155					160					
tca	acc	ata	gtc	aaa	aag	gaa	gga	ctt	ggt	gct	ctg	tgg	act	ggg	ctt	825
Ser	Thr	He	Val	Lys	Lys	Glu	Gly	Leu	Gly	Ala	Leu	Trp	Thr	Gly	Leu	
165					170					175					180	
ggt	cct	aat	att	gcc	cgc	aat	gct	att	ata	aat	gct	gct	gaa	ttg	gcc	873
Gly	Pro	Asn	He	Ala	Arg	Asn	Ala	He	He	Asn	Ala	Ala	Glu	Leu	Ala	
				185					190				•	195		
agt	tat	gat	caa	gtg	aaa	cag	atg	aag	tct	aga	atg	atg	gga	gat	tca	921
Ser	Tyr	Asp	GIn	Val	Lys	GIn	Met	Lys	Ser	Arg	Met	Met	Gly	Asp	Ser	
			200					205					210			
gcc	tac	aaa	agc	aca	ttt	gat	tgt	ttc	atc	aag	acg	ttg	aaa	aat	gat	969
Ala	Tyr	Lys	Ser	Thr	Phe	Asp	Cys	Phe	He	Lys	Thr	Leu	Lys	Asn	Asp	
		215					220					225				
ggg	cct	ctt	gct	ttt	tac	aag	ggg	ttt	atc	сса	aac	ttt	ggt	cgg	tta	1017
Gly	Pro	Leu	Ala	Phe	Tyr	Lys	Gly	Phe	Пe	Pro	Asn	Phe	Gly	Arg	Leu	
	230					235					240					
gga	tcg	tgg	aat	gtg	atc	atg	ttt	ttg	асс	ttg	gag	cag	gtc	aag	aag	1065
Gly	Ser	Trp	Asn	Val	He	Met	Phe	Leu	Thr	Leu	Glu	Gln	Val	Lys	Lys	
245					250					255					260	
ttc	ttc	atc	aaa	gag	gtg	cca	aat	taa	taca	ttg	aagt	cgga	ta g	gagt	agaaa	1119
Phe	Phe	He	Lys	Glu	Val	Pro	Asn									


265


aaaagggttt ttgtggaatt ttctctaccg gtgtggatcc tggcgagaga gaataaatct 1179 tectgactge teagatgttg tacettttt atgaatggtt ettttettat agaggacaga 1239 gaaaataaaa gaaaaattca ttgtcatgta ctctttttcc ccatttctgc tgagtagcag 1299 ctataccaag cagactitgt tgcttggctg ctgctaatct tgtagctgaa gaaaagtctt 1359 acattgagtg ataacgttgt tctctgcatc cattattttt cagagttact atttgacaca 1419 tgaaaagttt tiittiitt tiittitti aacaggcagc aaatagagga aicgaictca 1479 cgactatcct ctttattcat taacaggcat acaaacttag ggagagcatg cagggtatat 1539 atcaaaatat accettttat tagacatttt gegtacaeag ttggteetea aaegaetgta 1599 tctagcagcc aatttttaga ccacattaag acagagagaa acaagcagaa gaacagggta 1659 ccatacatac ataggtaata attaagatga tgaacatagc ataggttcat gatctacttc 1719 ttcttcacgt acacatgatg caccagctga atgggaatct tggtcaccat atggcatgaa 1779 agtacgtcat gtgcagacgt tatatagtgt tcttcttacc attcagcagc agcaccagag 1839 gcatcaaaca ctggggtctt gacagggtat gaggggtaca ttgcgatccc acacatgccg 1899 tagggagtca cattgcgttt gatttttatg tatccagcct gtccccacct agtgccccat 1959 gagtteetta caageeaata ateetteesa teeteegaac catateetat tateaceaca 2019 gcatggtcga tacgttgacc acatggtcca gcaaatacgc ccgaggtgta gtgttggaat 2079 ccagcgcccg aagcctcaag agcaacactg acaggttgct ttgcgactgc atactgtagg 2139 ctaacctcgt tgtacggaga aacattttca tacgcatcaa tcgaggtgac tttaataaga 2199 tttgctctgc aagttcctcg acgtcctgtg tacgggtaat ttttgaaaaa aggagaaggg 2259 cgggaagaag cgcgcgtctc tgctcgcgac gggttaattc ttcatatggc cacttcgagc 2319 atggcttcgg cagcctccag cttcgttctc actccggcct cccctccacc accacgccgg 2379 ttccccgtc cgccttcttt cccagggtcg caggaggctc gtggtggtgc gggccgagga 2439 gccaccgatc gggcccaaga gggggtcaaa ggttgatata tagttcttaa tttctttccc 2559 taaattttat tctcatctgt ggacgacctt ccatcggggt tttcgtccct ctcgcaggtg 2679 aagateteeg gaaggaatee taetggttea aeggtgtegg ateggtggtg getgttgate 2739 aggatccggc gactcgatac ccggtcgtgg ttcggttcac caaggtcaac tatgcgaacg 2799 tctcgaccaa caactacgca ctggacgaga tcctggaggt gaaatgaggg tcggcgggcg 2859
tggtcggtcg ggcatgtcac gatgatgtat tttcgcagtt ggtagtgtaa aataccatgt 2919
cattcgtgta aaactctttc gttcgccaaa tcctcagttg aaattttaat tcccagccag 2979
taaaaaaaaaa aa 2991

<210> 4 <211> 268 <212> PRT <213> Symplocarpus foetidus <400> 4 Met Gly Asp His Gly Pro Arg Thr Glu lle Ser Phe Ala Gly Ser Ser Arg Ala Ala Phe Ala Ala Cys Phe Ala Glu Leu Cys Thr lle Pro Leu Asp Thr Ala Lys Val Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly Asp Val Val Ala Leu Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala Thr lle Ala Arg Glu Glu Gly Leu Ser Ala Leu Trp Lys Gly lie Val Pro Gly Leu His Arg Gln Cys Leu Phe Gly Gly Leu Arg Ile Gly Leu Tyr Glu Pro Val Lys Ser Phe Tyr Val Gly Asp Asn Phe Val Gly Asp lle Pro Leu Ser Lys Lys lle Leu Ala Gly Leu Thr Thr Gly Ala Leu Ala lle lle Val Ala Asn Pro Thr Asp Leu Val Lys Val Arg Leu Gln 

Ser	Glu	Gly	Lys	Leu	Pro	Pro	Gly	Val	Pro	Arg	Arg	Tyr	Ser	Gly	Ala
145					150					155					160
Leu	Asn	Ala	Tyr	Ser	Thr	He	Val	Lys	Lys	Glu	Gly	Leu	Gly	Ala	Leu
				165					170					175	
Trp	Thr	Gly	Leu	Gly	Pro	Asn	He	Ala	Arg	Asn	Ala	He	ile	Asn	Ala
			180					185					190		
Ala	Glu	Leu	Ala	Ser	Tyr	Asp	Gln	Val	Lys	Gln	Met	Lys	Ser	Arg	Met
		195					200					205			
Met	Gly	Asp	Ser	Ala	Tyr	Lys	Ser	Thr	Phe	Asp	Cys	Phe	He	Lys	Thr
	210					215					220				
Leu	Lys	Asn	Asp	Gly	Pro	Leu	Ala	Phe	Tyr	Lys	Gly	Phe	He	Pro	Asn
225					230					235					240
Phe	Gly	Arg	Leu	Gly	Ser	Trp	Asn	Val	He	Met	Phe	Leu	Thr	Leu	Glu
				245					250					255	
GIn	Val	Lys	Lys	Phe	Phe	He	Lys	Glu	Val	Pro	Asn				
			260					265							

Fig.1





### Fig.3 Top

**-**

ifuceB stuce terumen ucel numen uce2 numen uce2	1MGDHGPRTEISFAGSSR-AAFAACFAEICTTPLDTAKVRUQLQKKAVTGDV-VALFNI 1 MGGGDHGGKSDISFAGIFASSAFAACFAEACTTPLDTAKVRLQLQKKAVEGCG-LALPKY 1MVAAGKSDLSLPKTFACSAFAACVGEVCTTPLDTAKVRLQLQKSAFTLAGDVTLPKY 1MVAGKSDLSLPKTFACSAFAACVGEVCTTPLDTAKVRLQLQKSAFTLAGDVTLPKY 1MGGLTASDVHPTLGVQLFSAPIAACLADVITHPLDTAKVRLQUGGECPTSSVIRY 1	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
	YNO TAIL MICHAEL TO TO TO A TO TO AND A SEC A SE	אַ
fucea	1MGDRGPRIEISFAGSSR-AAFAACFAELCTIPLLIAKVALQLQKAAVIGDV-VALFAI	י ע
sfuces stuce ttrumen numan UCP1	1MGDHGFRTEISFAGSSR-AAFAACFAELCT FLDIAKVHUQLQKKAVIGDV-VALFAL 1 MGGGDHGGKSDISFAGIFASSAFAACFAEACTTPLDTAKVRLDLQKKAVEGDG-LALPKY 1MVAAGKSDLSLPKTFACSAFAACVGEVCTIPLDTAKVRLDLQKSAFTLAGDVTLPKY 1MGGLTASDVHPTLGVQLFSAPIAACLADVITHPLDTAKVRLDVQGECPTSSVIRY	5 2 3 6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
uman UCP2	1 -MVGFKATDVPPTATVKFLGAGTAACIADLITHPLDTAKVRLDIQGESQGPVRATASAQY 1 -MVGLKPSDVPPTMAVKFLGAGTAACFADLVTHPLDTAKVRLDIQGENQ-AVQTARLVQY	2 S

**ゴ** ゴ

117 RGVMGTILTMVRTECPRSLYNGLVAGLQRQMSFASVRIGLYDSVKQFYT----KGSZHAS RGVLGTILTMVRTEGPCSPYNGLVAGLQRQNSFASIRIGLYDSVKQVYTP---KGADNSS KGVLGTITAVVKTEGRÆKLYSGLPAGLQRQISSASLRIGLYDTVQEFLTA---GKETAPS RCMLGTVATIAREEGLSALWKGIVPGLHRQCLFGGLRIGLYEPVKSFYVG--DNFVGDIP RGMIGIVATIAREEGISALHKGIVPGLHRQCLFGGLRIGLYEPVKSFYVG--DNFVGDIP RGLLGTVGTIAREEGLRSLHKGVVPGLHRQCLFGGLRIGMYEPVKNLYVFTGKDFVGDVP RGLLGTVGTLAKEEGLASLWKGIVPGLHRQCIYGGLRIGHYEPVKNLYVG--KDHVGDVP 9 58 56 9 UCP3 UCP2 human UCP1 SfUCPB AtPUMP SEUCPA human human Stucp

# Fig.3 Middle

### III

		TEMILED VAN TADDAGGE, Dag military	1 7 1
SfUCPA	115		
Sfuceb	115		171
Stuce	118		174
AtPUMP	118		176
human UCP1	113		168
human UCP2	116	IGSRLLAGSTTGALAVAVA¢PTDVVKVRFDAQARAGGGRRYQSTVNAYKTIAREE	170
human UCP3	116	LTTRILAGCTTGAMAVTCAQPTDVVKVREDASIHLGPSRSDRKYSGTMDAYRTIAREE	173
		* .* .*	
		Λ	
Sfucea	172	172 GLGALWTGLGPNIARNAIINNAELASYDQVKQTILKLPGFSDNIFTHILAGLGAGFFA 229	223
Sfucrb	172	172 GLGALMTGLGPNIARNAIINAAELASYDQVKQ	203
Stuce	175		232
AtPUMP	177	177 GVRALWTVLGPNVARNAIINAAELASYDQVKETILKIPGFTDNVVTHILSGLFTGAGFFA	236
human UCP1	169	169 GLIGLWRGTTPNLMRSVIINCTELVTYDLMKEAFVKNNILADDVPCHLVSALLAGFCA 226	226
human UCP2	171	171 GFRGLWKGISPNVARNALVNCAELVTYDLIKDALLKANLMIDDLPCHFISAFGAGFCT 228	277
human UCP3	174	174 GVRGLWKGTLPNIMRNAIVNCAEVVTYDILKEKLLDYHLLTDNFPCBFVSAFGAGFCA 233	77

## Fig.3 Down

285	250	288	295	284	286	589										
230 VCIGGPVDVARSRANGDSAYRSTFDCFIRTLKNDGLLAFYRGFIPNFGRLGSWNVI	hksramedsaiksifdcfikilkndgplafykgfipnfgrlgswnvi	VCIG4PVDVVKSRMMGDSAYKNTLDCFVKTLKNDGPLAFYKGFIPNFGRLGSWNVI 2	VCIC4PVDVVKSRAMGDSG-AYKGTIDCFVKTLKSDGPHAFYKGFIPNFGRLGSFTWNVI ;	TAMS & PVDVVKTRA INSPPCQYKSVPNCAMKVFTNEGPTAFFKGLVPSFLALGSWNVI	TVIASPVDVVKTRHMSALGQYSSAGECALIMQKEGPRAFYKGFMPSFIRLGSWNVV	TVVA PVDVVKTRHANSPPGQYFSPLDCMIKHVAQEGPTAFYKGFTPSFLRLGSWNVV			- 303	- 268	- 306	- 316	r 307	309	F 312	
MGDSAYP	MGDSAY	MGDSAYR	MGDSG-AYP	INSPPGQYF	MSALGQYS	MSPPGQYE	•		EVPN	EVPN	LESP	QAKKYVRELDASKRN	QLKRELSKSRQTMDCAT	QLKRALMAACTSREAPF	QLKRALMKVQMLAESPF	
PVDVMRSRM	MKSRW	PVDVVKSRM	PVDVVKSRM	PVDVVKTRE	PVDVVKTRK	PVDVVKTRH	*.	PNBD	286 MFLILEQVKKFFIKEVPN	HFLILEQVRKFFIKEVPN	HFLTTEQAKKFVKSLESP	<b>EQAKKYVRE</b>	EQLKRELSR	EQLKRALMA	EQLKRALMR	
VCIG	1	VCICS							MFLIT	MILIT	MFLTL	MFLTLE	MEVCFE	MEVIYE	MEVITE	*
 230	204	233	237	227	229	232			286	251	289	296	285	289	290	
SfUCPA	SfUCPB	Stuce	AtPUMP	human UCP1	human UCP2	human UCP3			SfUCPA	SfUCPB	Stuce	AtPUMP	human UCP1	human UCP2	human UCP3	

Fig.4





Fig.5

### Intermembrane space



Mitochondrial matrix

Fig.6






Fig.7

#### Intermembrane space



Mitochondrial matrix

Fig.8



Effective March 1998

### DECKARATION AND POWER OF ATTORNEY FOR U.S. PATENT APPLICATION

(X) Original () Supplemental () Substitute (X) PCT () DESIGN

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that I verily believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Title:	PLANT THERMOGENIC	<b>GENES</b>	AND	<b>PROTEINS</b>

Litera Marian
which is described and claimed in:
the attached specification, or
the specification in application Scrial No, filed December 14, 2001, and with amendments through, or
the specification in International Application No. <u>PCT/JP00/03806</u> , filed <u>June 12, 2000</u> , and as amended on

I hereby state that I have reviewed and understand the content of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge my duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim priority benefits under Title 35, United States Code, §119 (and §172 if this application is for a Design) of any application(s) for patent or inventor's certificate listed below and have also identified below any application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

COUNTRY	APPLICATION NO.	DATE OF FILING	PRIORITY CLAIMED
Japan	1999-167439	June 14, 1999	Yes

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I scknowledge the duty to disclose information material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	U.S. FILING DATE	STATUS: PATENTED, PENDING, ABANDONED	

And I hereby appoint Michael R. Davis, Reg. No. 25,134; Matthew M. Jacob, Reg. No. 25,154; Warren M. Cheek, Jr., Reg. No. 33,367; Nils Pedersen, Reg. No. 33,145; Charles R. Watts, Reg. No. 33,142; and Michael S. Huppert, Reg. No. 40,268, who together constitute the firm of WENDEROTH, LIND & PONACK, L.L.P., as well as any other attorneys and agents associated with Customer No. 000513, to prosecute this application and to transact all business in the U.S. Patent and Trademark Office connected therewith.

I hereby authorize the U.S. attorneys and agents named herein to accept and follow instructions from NISHIZAWA & ASSOCIATES as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and myself. In the event of a change in the persons from whom instructions may be taken, the U.S. attorneys named herein will be so notified by me.

Direct Correspondence to Customer No:



PATENT TRADEMARK OFFICE

Direct Telephone Calls to:

WENDEROTH LIND & PONACK LL.P. 2033 "K" Street, N.W., Suite 800 Washington, D.C. 20006-1021

Phone:(202) 721-8200 Fax:(202) 721-8250

1-00

Full Name of First Inventor	FAMILY NAME ITO	first civen name Kikukatsu	JECOND GIVEN NAME
Residence & Citizenship	Iwate	STATE OR COUNTBY  Japan JP	сочитку от сптижизите Јарап
Post Office Address	Estate Mide Japan	origa-oka 207, 17-7, Mide	STATE OR COUNTRY ZIP CODE Origa-oka 1-chome, Morioka-shi, Iwate,
Full Name of Second Inventor	FAMILY NAME	First civen name	SECOND GIVEN NAME
Residence & Citizenship	СПҮ	STATE OR COUNTRY	COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS	CITY	STATE OR COUNTRY ZIP CODE
Full Name of Third Inventor	Family name	first civen name	second given name
Residence & Citizenship	CITY	STATE OR COUNTRY	COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS	CITY	STATE OR COUNTRY ZIP CODE
Full Name of Fourth Inventor	Family name	FIRST GIVEN NAME	SECOND GIVEN NAME
Residence & Citizenship	CITY	STATE OR COUNTRY	COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS	CITY	STATE OR COUNTRY ZIP CODE
Full Name of Fifth Inventor	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
Residence & Citizenship	aтy	STATE OR COUNTRY	COUNTRY OF CTITZENSHIP
Post Office Address	ADDRESS	СПҮ	STATE OR COUNTRY ZIF CODE
Full Name of Sixth Inventor	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
Residence & Citizenship	CITY	STATE OF COUNTRY	COUNTRY OF CITEZENSHIP

		COTO .	T. T. O. COLDINA	ZIP CODE
Post Office	ADDRESS	cmy	STATE OR COUNTRY	ZF COBE
Address				

I further declare that all statements made herein of my own knowledge are true, and that all statements on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1st Inventor Kikukatsu ITO	Date	January 11, 2002				
Kikukatsu ITO		``				
2nd Inventor	Datc _					
3rd Inventor	Date					
4th Inventor	Date					
5th Inventor	Date					
6th Inventor	Date					
The above application may be more particularly identified as f	ollows:					
U.S. Application Serial No Filing I	Date December 14, 2001					
Applicant Reference Number 99-F-071US/YS Atty Docket No. 2001 1838A						

Title of Invention PLANT THERMOGENIC GENES AND PROTEINS